

Evidence-Based Nursing

Contents

| | | | |
|---------------------------------------------------------------------------------------------------------------------------------------------------|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|
| Purpose and procedure | 98 | Oestrogen plus progestin increased venous thromboembolic disease in postmenopausal women with coronary artery disease | 123 |
| EBN users' guide | | Review: both glucosamine and chondroitin are effective for osteoarthritis, but the magnitude of effect is unclear | 124 |
| Evaluation of studies of treatment or prevention interventions | 100 | An advance directive programme in nursing homes reduced health services use without affecting patient satisfaction | 125 |
| Implementation forum | | Prompt hospital discharge with home care improved physical health and community reintegration and reduced initial length of hospital stay after acute stroke | 126 |
| Developing organisational systems and culture to support evidence-based practice: the experience of the Evidence-Based Ward Project | 103 | Early discharge plus home based rehabilitation reduced length of initial hospital stay but did not improve health related quality of life in patients with acute stroke | 127 |
| Acknowledgments | 105 | Causation | |
| Other articles noted | 106 | Restraint use and rear seating were associated with fewer serious injuries and deaths for children in motor vehicle crashes | 128 |
| Treatment | | Siblings of children who had unintentional injuries were at increased risk of injury for 90 days after the event | 129 |
| Review: daily maintenance inhaled corticosteroids are not effective for children with episodic viral wheeze | 113 | Dietary patterns including fruits, vegetables, whole grains, low fat dairy, and lean meats lowered mortality risk in women | 130 |
| Supplemental oxygen did not reduce progression of pre-threshold retinopathy of prematurity in infants | 114 | Qualitative | |
| Nurse home visits during pregnancy and early childhood had positive effects on aspects of maternal life course 3 years later | 115 | Perceived breast milk inadequacy was related to sociocultural influences and the behaviour of women and their babies | 131 |
| Cognitive behavioural therapy improved psychiatric symptoms in drug refractory schizophrenia | 116 | Homeless people preserved former identities, devalued current identities, and envisioned a non-homeless future | 132 |
| Automated telephone assessment and education with nurse follow up improved self care and glycaemic control in patients with diabetes | 117 | Dealing with a family member who has a mental illness was a long term, frustrating, and confusing process before acceptance occurred | 133 |
| Review: aromatherapy massage is associated with small, transient reductions in anxiety | 118 | Nurses and family caregivers of elderly relatives engaged in 4 evolving types of relationships | 134 |
| A 0.5% chlorhexidine gluconate in 70% isopropyl alcohol swab was more effective than 2 other methods for intravenous skin antisepsis | 119 | Helping, mutual sharing, committing, and benefiting described the peer advisor experience of providing social support | 135 |
| Immune enhancing enteral nutrition reduced mortality and acquired infections in intensive care unit patients with sepsis | 120 | Nursing home residents used 6 strategies to manage urinary incontinence | 136 |
| Daily interruption of sedative infusions reduced duration of mechanical ventilation and intensive care unit stay in critically ill patients | 121 | Glossary | Inside back cover |
| Pelvic floor re-education reduced incontinence 1 year after radical prostatectomy | 122 | | |